

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

EP 0 610 049 B1

11

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
03.11.1999 Bulletin 1999/44

(51) Int Cl.<sup>6</sup>: C12P 41/00, C12P 7/42  
// C12P7:42,(C12R1/05, 1:13,  
1:38)

(21) Application number: 94300705.4

(22) Date of filing: 31.01.1994

(54) **Process for producing optically active alpha-hydroxycarboxylic acid having phenyl group**

Verfahren zur Herstellung von Phenylgruppe enthaltende optisch-aktiven  
Alpha-Hydroxycarbonsäuren

Procédé de préparation d'acides alphahydroxy carboxyliques optiquement actifs contenant un groupe  
phénylique

(84) Designated Contracting States:  
DE FR GB

(30) Priority: 03.02.1993 JP 3727693  
08.09.1993 JP 24602893

(43) Date of publication of application:  
10.08.1994 Bulletin 1994/32

(73) Proprietor: MITSUBISHI RAYON CO., LTD.  
Minato-ku, Tokyo (JP)

(72) Inventors:  
• Hashimoto, Yoshihiro,  
c/o Nitto Chem. Ind. Co. Ltd.  
Tsurumi-ku, Yokohama-shi, Kanagawa (JP)  
• Endo, Takakazu, c/o Nitto Chem. Ind. Co. Ltd.  
Tsurumi-ku, Yokohama-shi, Kanagawa (JP)

- Tamura, Koji, c/o Nitto Chem. Ind. Co. Ltd.  
Tsurumi-ku, Yokohama-shi, Kanagawa (JP)
- Hirata, Yuji, c/o Nitto Chem. Ind. Co. Ltd.  
Tsurumi-ku, Yokohama-shi, Kanagawa (JP)
- Kobayashi, Etsuko, c/o Nitto Chem. Ind. Co. Ltd.  
Tsurumi-ku, Yokohama-shi, Kanagawa (JP)

(74) Representative: Pearce, Anthony Richmond  
MARKS & CLERK,  
Alpha Tower,  
Suffolk Street Queensway  
Birmingham B1 1TT (GB)

(56) References cited:  
EP-A- 0 348 901 EP-A- 0 449 648  
EP-A- 0 486 289

EP 0 610 049 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

## Description

## FIELD OF THE INVENTION

5 [0001] This invention relates to a biological process for producing an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group. More particularly, it relates to a process for producing an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group as represented by formula (I) shown below, by the action of a microorganism capable of asymmetrically hydrolyzing a nitrile group of a racemic  $\alpha$ -hydroxynitrile represented by formula (II) shown below. The optically active  $\alpha$ -hydroxycarboxylic acid of formula (I) is of industrial importance as a starting material for synthesizing pharmaceuticals and agrochemicals, such as antibiotics, drugs acting on the sympathetic nervous system, anti-diabetic agents, and as a resolving reagent.

## BACKGROUND OF THE INVENTION

15 [0002] Known processes for producing an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group include optical resolution of racemates by crystallization or chromatography and asymmetric synthesis through organochemical procedures. These processes generally involve complicated operations yet give only a low yield of a product.


[0003] To overcome these problems, biological processes utilizing microorganisms have been proposed. For example, it has been proposed to asymmetrically hydrolyze substituted or unsubstituted mandelonitrile or substituted or unsubstituted mandelamide by the action of a microorganism belonging to the genus Alcaligenes, Pseudomonas, Rhodopseudomonas, Corynebacterium, Acinetobacter, Bacillus, Mycobacterium, Rhodococcus or Candida to obtain optically active mandelic acid or a derivative thereof, as disclosed in European Patent Publication No. 0 384 901 A (corresponding to JP-A-2-84198 (the term "JP-A" as used herein means an "unexamined published Japanese patent application")) and Applied and Environmental Microbiology, Vol. 57, pp. 3028-3032 (1991). It is also known to obtain a predominant proportion of R(-)-mandelic acid or a derivative thereof directly from racemic mandelonitrile or a derivative thereof by the action of a microorganism belonging to the genus Pseudomonas, Alcaligenes, Acinetobacter, Caseobacter, Nocardia, Bacillus, Brevibacterium or Aureobacterium as disclosed in U.S. Patent 5,223,416 (corresponding to JP-A-4-218385, JP-A-4-99495 and JP-A-4-99496).

25 [0004] However, the above publications do not disclose producing an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group represented by formula (I) nor what activity the above microorganisms have relating to producing an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group represented by formula (I).

## SUMMARY OF THE INVENTION


35 [0005] The present inventors searched extensively for a microorganism capable of producing an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group with industrial advantages. As a result, for such purposes, they found it effective to use a microorganism belonging to the genus Rhodococcus, Alcaligenes, Brevibacterium or Pseudomonas.

[0006] The present invention relates to a process for predominantly producing an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group, as represented by formula (I):



wherein X represents a hydrogen atom, a hydroxyl group, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a thioalkyl group, a halogen atom, a phenyl group, a phenoxy group, an amino group or a nitro group, which is bonded to the o-, m-, or p-position; and n represents 1 or 2,

50 which comprises reacting a microorganism capable of asymmetrically hydrolyzing a nitrile group of a racemic  $\alpha$ -hydroxynitrile represented by formula (II):



wherein X and n are as defined above,

and wherein said microorganism belongs to the genus Rhodococcus, Alcaligenes, Brevibacterium or Pseudomonas or treated cells thereof with a racemic  $\alpha$ -hydroxynitrile represented by formula (II) or a mixture of an aldehyde corresponding to said nitrile and prussic acid in a neutral to basic aqueous medium.

- 5 [0007] The present invention is based on the fact that the  $\alpha$ -hydroxynitrile of formula (II) easily racemizes in a neutral to basic aqueous medium in the dissociation equilibrium between the  $\alpha$ -hydroxynitrile and the corresponding aldehyde and prussic acid. The inventors found that the  $\alpha$ -hydroxynitrile of formula (II) can be directly converted to an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group represented by formula (I) with predominance over a D-form or an L-form by coupling the above-mentioned racemization reaction with the above-mentioned microorganism. The term "with predominance" as used herein means that a D-form or an L-form is obtained from a racemic compound in a yield of from 50 to 100% based on the amount of reactant according to formula (II).

#### DETAILED DESCRIPTION OF THE INVENTION

- 15 [0008] The microorganism which can be used in the present invention belongs to the genus Rhodococcus, Alcaligenes, Brevibacterium or Pseudomonas and is capable of asymmetrically hydrolyzing a nitrile group of a racemic  $\alpha$ -hydroxynitrile of formula (II) to produce and accumulate an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group, represented by formula (I), in high concentration.

- 20 [0009] The microorganism typically includes Rhodococcus sp. HT29-7 (FERM BP-3857), Alcaligenes sp. BC35-2 (FERM BP-3318), Alcaligenes sp. BC12-2 (FERM P-11263), Alcaligenes sp. BC20 (FERM P-11264), Alcaligenes sp. BC24 (FERM P-12063), Pseudomonas sp. BC-18 (FERM BP-4536), and Brevibacterium acetylicum (IAM 1790). Of these microorganisms, Rhodococcus sp. HT29-7, Alcaligenes sp. BC35-2, BC12-2, BC20 and BC24, and Pseudomonas sp. BC-18 are novel strains which the inventors isolated from soil and deposited with National Institute of Bioscience & Human Technology (formerly, Fermentation Research Institute), Agency of Industrial Science & Technology under the respective deposit receipt numbers. Morphological and physiological properties of these strains are described below. Brevibacterium acetylicum IAM 1790 is a known strain and available at Institute of Applied Microbiology, The University of Tokyo (IAM).

30	HT29-7:	
	Shape	polymorphic bacillus
	Gram's stain	+
	Spore	-
	Motility	-
35	Color of colony	pink to orange
	Rod-coccus cycle	+
	Extension of peripheral cells of colony	observed
	Formation of aerial hypha	not observed
40	Oxidase	-
	Catalase	+
	Behavior toward oxygen	aerobic
	Diamino acid of cell wall	meso-diaminopimelic acid
	Glycolyl test	+(glycolyl type)
45	Sugar composition of cell wall	
	Arabinose	+
	Galactose	+
	Existence of quinone	MK-9 (H <sub>2</sub> )

50	BC35-2, BC12-2, BC20, and BC24:	
	Shape	bacillus
	Gram's stain	-
55	Spore	-
	Motility	+
	Flagellum	peripheral

(continued)

## BC35-2, BC12-2, BC20, and BC24:

Oxidase	+
Catalase	+
OF test	alkalization
Production of 3-ketolactose	-
Existence of quinone	Q-8

## BC-18:

Shape	bacillus
Gram's stain	-
Spore	-
Motility	+
Flagellum	polar
Oxidase	+
Catalase	+
OF test	0

[0010] The above-described taxonomical properties were examined by referring to Bergey's Manual of Systematic Bacteriology (1986). The HT29-7 strain belongs to the genus Rhodococcus; BC35-2, BC12-2, BC20 and BC24 strains belong to the genus Alcaligenes; and the BC-18 strain belongs to the genus Pseudomonas.

[0011] Typical examples of the compounds of formula (II) which can be used as a substrate in the present invention include phenyllactonitrile, 4-phenyl- $\alpha$ -hydroxybutyronitrile, 3-(2-methoxyphenyl)-lactonitrile, 3-(3-methoxyphenyl)-lactonitrile, 3-(4-methoxyphenyl)-lactonitrile, 4-(4-fluorophenyl)- $\alpha$ -hydroxybutyronitrile, 4-(2-chlorophenyl)- $\alpha$ -hydroxybutyronitrile, 4-(4-bromophenyl)- $\alpha$ -hydroxybutyronitrile, 4-(2-trifluoromethylphenyl)- $\alpha$ -hydroxybutyronitrile, 4-(3-trifluoromethylphenyl)- $\alpha$ -hydroxybutyronitrile and 4-(2-hydroxyphenyl)- $\alpha$ -hydroxybutyronitrile.

[0012] Cultivation of the microorganism is carried out by using usual media containing assimilable carbon sources (e.g., glycerol, glucose, saccharose, malt extract, lactose, and fructose); assimilable nitrogen sources (e.g., casamino acid, meat extract, and yeast extract); and inorganic nutrients essential for growth (magnesium chloride, sodium sulfate, calcium chloride, manganese sulfate, iron chloride, and zinc sulfate).

[0013] To obtain increased enzyme activity, an enzyme inducer preferably is added to the culture medium in the initial or middle stage of cultivation, in such a concentration as not to significantly inhibit growth. Suitable enzyme inducers include nitriles (e.g., cinnamonnitrile, benzyl cyanide, isobutyronitrile,  $\beta$ -phenylpropionitrile, benzonitrile, 2-, 3- or 4-cyanopyridine, 1-cyclohexenylacetonitrile,  $\epsilon$ -caprolactam,  $\gamma$ -butyronitrile, and o-aminobenzonitrile); and amides (e.g., isobutylamide, phenylacetamide, and 4-pyridinecarboxylic acid amide).

[0014] Culturing is aerobically conducted at a pH of 4 to 10 and a temperature of 5 to 50°C, for a period of about 1 to 7 days, until the maximum activity is reached.

[0015] The asymmetric hydrolysis reaction can be carried out by suspending microbial cells harvested from the culture or treated microbial cells (e.g., dried cells, ruptured cells, a crude or purified isolated enzyme, immobilized microbial cells or an immobilized enzyme) in an aqueous medium (e.g., water and a buffer solution), and bringing a racemic  $\alpha$ -hydroxynitrile of formula (II) or a mixture of an aldehyde corresponding to the nitrile and prussic acid into contact with the cell suspension. The reaction system should be maintained near neutral or basic in order to racemize the  $\alpha$ -hydroxynitrile of formula (II). That is, the pH of the reaction system should be kept within a range of 6 to 11, and preferably 7 to 10.

[0016] The concentration of the substrate in the reaction system usually ranges from 0.1 to 10% by weight, and preferably from 0.2 to 5.0% by weight in terms of the racemic  $\alpha$ -hydroxynitrile of formula (II), while varying depending on sensitivity of the enzyme to the aldehyde corresponding to the racemic  $\alpha$ -hydroxynitrile of formula (II) or prussic acid.

[0017] To reduce enzyme denaturation by an aldehyde, sodium sulfite, acid sodium sulfite, sodium dithionite, potassium sulfite, acid potassium sulfite, potassium dithionite, ammonium sulfite, acid ammonium sulfite, or ammonium dithionite in an amount of 1 to 1000 mM may be added.

[0018] The microorganism usually is used in an amount of 0.01 to 5.0% by weight on a dry basis based on the substrate. The reaction usually is conducted at a temperature of 0 to 50°C, and preferably 10 to 30°C, for a period of 0.1 to 100 hours.

[0019] The reaction product, i.e., an optically active  $\alpha$ -hydroxycarboxylic acid of formula (I), can be isolated from the reaction mixture by known procedures. For example, the microbial cells may be removed by centrifugation, and if desired, granular components, proteins, and polysaccharides may be removed by ultrafiltration or the like. The supernatant may be treated with activated carbon. Then, the supernatant is concentrated under reduced pressure or extracted with an organic solvent in an acidic condition, followed by repeated recrystallization from benzene, etc. to obtain a high purity crystal.

[0020] The present invention provides an industrially excellent process for preparing an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group represented by formula (I) at almost quantitative selectivity and at high optical purity.

[0021] The present invention will now be illustrated in greater detail with reference to Examples, but it should be understood that the present invention is not deemed to be limited thereto. All percentages are by weight unless otherwise indicated.

#### EXAMPLE 1

##### Preparation of L-3-Phenyllactic Acid

##### (1) Culturing:

[0022] *Rhodococcus* sp. HT29-7 (FERM BP-3857), *Alcaligenes* sp. BC35-2 (FERM BP-3318) or *Brevibacterium acetylicum* IAM 1790 was aerobically cultured in a medium having the following composition, with 0.05% benzyl cyanide added as an inducer, at 30°C for 72 hours.

Medium Composition:	
Glycerol	20 g
Yeast extract	3 g
Potassium primary phosphate	6.8 g
Sodium secondary phosphate	7.1 g
Sodium sulfate	2.8 g
Magnesium chloride	0.4 g
Calcium chloride	0.04 g
Manganese sulfate	0.03 g
Iron chloride	0.006 g
Zinc sulfate	0.003 g
Distilled water	1000 ml
pH = 7.5	

##### (2) Asymmetric Hydrolysis:

[0023] Microbial cells harvested from the culture were washed with a 50 mM phosphoric acid buffer solution (pH=8.2) by centrifugation and suspended in the same buffer solution containing 10 mM phenyllactonitrile at such a cell concentration as to have an optical density at 630 nm ( $OD_{630}$ ) of 20. The cell suspension was allowed to react at 30°C for 96 hours while shaking.

[0024] After completion of the reaction, the microbial cells were removed by centrifugation. The content of phenyllactic acid in the supernatant was determined by liquid chromatography (column: Wakosil® ODS 5C18; carrier solution: 0.1M phosphoric acid:acetonitrile=3:1 by volume; monitor: 254 nm). The supernatant was adjusted to pH 12 with 6N NaOH and extracted twice with an equal amount of ethyl acetate to remove any unreacted phenyllactonitrile. The aqueous layer was adjusted to pH 1.2 with sulfuric acid and extracted twice with an equal amount of ethyl acetate. The extract was evaporated to dryness in an evaporator, and the residue was dissolved in water and analyzed on an optical resolution column (MCI gel CRS-10W; carrier solution: 2mM  $CuSO_4 \cdot 5H_2O$ :acetonitrile=85:15 by volume). The results obtained are shown in Table 1 below.

TABLE 1

Microorganism	Yield	Optical Purity (L-Form)
	(mM)	(%ee)
HT29-7	6.0 (60%)	60
BC35-2	0.55 (5.5%)	45
IAM 1790	9.9 (99%)	10

**EXAMPLE 2**Preparation of L-3-Phenyllactic Acid

[0025] Microbial cells of HT29-7 strain or IAM 1790 strain harvested in the same manner as in Example 1 were suspended in a phosphoric acid buffer solution (pH=8.2) containing 10 mM phenylaldehyde and 10 mM potassium cyanide to a cell concentration of  $OD_{630} = 20$ . The cell suspension was allowed to react at 30°C for 96 hours with shaking.

[0026] The reaction mixture was worked-up in the same manner as in Example 1. The phenyllactic acid content and its optical purity were determined in the same manner as in Example 1, with the results shown in Table 2 below.

TABLE 2

Microorganism	Yield	Optical Purity (L-Form)
	(mM)	(%ee)
HT29-7	7.1 (71%)	64
IAM 1790	9.8 (98%)	14

**EXAMPLE 3**Preparation of L-4-Phenyl- $\alpha$ -hydroxybutyric Acid

[0027] Microbial cells of HT29-7 strain harvested in the same manner as in Example 1 were suspended in a phosphoric acid buffer solution (pH=8.2) containing 10 mM 4-phenyl- $\alpha$ -hydroxybutyronitrile to a concentration of  $OD_{630} = 20$ . The cell suspension was allowed to react at 30°C for 96 hours with shaking. Further, the same reaction was carried out in the presence of 100 mM sodium sulfite for reducing enzyme denaturation by the aldehyde.

[0028] The reaction mixture was worked-up in the same manner as in Example 1. The 4-phenyl- $\alpha$ -hydroxybutyric acid content and the optical purity were determined in the same manner as in Example 1, with results shown in Table 3 below.

TABLE 3

Microorganism	Sodium Sulfite	Yield	Optical Purity (L-Form)
		(mM)	(%ee)
HT29-7	not added	9.4 (94%)	87
"	added	9.9 (99%)	91

**EXAMPLE 4**Preparation of L-3-Phenyllactic Acid

## (1) Culturing:

[0029] *Pseudomonas* sp. BC-18 (FERM BP-4536) was aerobically cultured in a medium having the following composition, with 0.03% 1-cyclohexenylacetonitrile added as an inducer, at 30°C for 3 days.



## Medium Composition:

Glycerol	20 g
Yeast extract	6 g
Metal salt mixed solution*	5 ml
1M Sodium sulfate	2 ml
50 mM Phosphoric acid buffer solution (pH=7.5)	993 ml

\* Sodium sulfate (56 g), magnesium chloride (8 g), calcium chloride (0.8 g), manganese sulfate (0.6 g), iron chloride (0.12 g) and zinc sulfate (0.06 g) in distilled water (100 ml)

## (2) Asymmetric Hydrolysis:

[0030] Microbial cells harvested from the culture were washed with a 50 mM phosphoric acid buffer solution (pH=7.5) by centrifugation and suspended in a phosphoric acid buffer solution having a prescribed pH and containing 20 mM phenyllactonitrile at a cell concentration of  $OD_{630}=29.0$ . The cell suspension was allowed to react at 30°C for 24 hours while shaking.


[0031] After completion of the reaction, the microbial cells were removed by centrifugation. The content of phenyllactic acid in the supernatant was determined by liquid chromatography (column: Wakosil ODS 5C18; carrier solution: 0.1M phosphoric acid:acetonitrile=7:3 by volume; monitor: 254 nm). The supernatant was adjusted to pH 9.0 with 6N NaOH and extracted twice with an equal amount of ethyl acetate to remove any unreacted phenyllactonitrile. The aqueous layer was adjusted to pH 2.0 with sulfuric acid and extracted twice with an equal amount of ethyl acetate. The extract was evaporated to dryness in an evaporator, and the residue was dissolved in water and analyzed on an optical resolution column (MCI gel CRS-10W; carrier solution: 2mM  $CuSO_4 \cdot 5H_2O$ :acetonitrile=85:15 by volume). The results obtained are shown in Table 4 below.

TABLE 4


Microorganism	pH	Yield	Optical Purity (L-Form)
		(mM)	(%ee)
BC-18	6.0	12.37 (61.9%)	21.1
"	6.5	11.35 (56.8%)	42.5
"	7.0	11.15 (55.8%)	48.0
"	7.5	11.00 (55.0%)	63.8
"	8.0	11.02 (55.1%)	61.4
"	8.5	14.05 (70.3%)	74.1
"	9.0	13.90 (69.5%)	73.8

## Claims

1. A process for predominantly producing an optically active  $\alpha$ -hydroxycarboxylic acid having a phenyl group as represented by formula (I):



wherein X represents a hydrogen atom, a hydroxyl group, an alkyl group having from 1 to 3 carbon atoms, an alkoxy group having from 1 to 3 carbon atoms, a thioalkyl group, a halogen atom, a phenyl group, a phenoxy group, an amino group or a nitro group, which is bonded to the o-, m-, or p-position; and n represents 1 or 2, which comprises reacting a microorganism capable of asymmetrically hydrolyzing a nitrile group of a racemic  $\alpha$ -hydroxynitrile represented by formula (II):




wherein X and n are as defined above,

and wherein the microorganism belongs to the genus Rhodococcus, Alcaligenes, Brevibacterium or Pseudomonas or treated cells thereof with a racemic  $\alpha$ -hydroxynitrile represented by formula (II) or a mixture of an aldehyde corresponding to said nitrile and prussic acid in a neutral to basic aqueous medium.

2. A process as claimed in Claim 1, wherein said microorganism has the properties of Rhodococcus sp. HT29-7 (FERM BP-3857), Alcaligenes sp. BC35-2 (FERM BP-3318), Alcaligenes sp. BC12-2 (FERM P-11263), Alcaligenes sp. BC20 (FERM P-11264), Alcaligenes sp. BC24 (FERM P-12063), Pseudomonas sp. BC-18 (FERM BP-4536), or Brevibacterium acetylicum (IAM 1790).
3. The process as claimed in Claim 1, wherein  $\alpha$ -hydroxynitrile is phenyllactonitrile, 4-phenyl- $\alpha$ -hydroxybutyronitrile, 3-(2-methoxyphenyl)-lactonitrile, 3-(3-methoxyphenyl)-lactonitrile, 3-(4-methoxyphenyl)-lactonitrile, 4-(4-fluorophenyl)- $\alpha$ -hydroxybutyronitrile, 4-(2-chlorophenyl)- $\alpha$ -hydroxybutyronitrile, 4-(4-bromophenyl)- $\alpha$ -hydroxybutyronitrile, 4-(2-trifluoromethylphenyl)- $\alpha$ -hydroxybutyronitrile, 4-(3-trifluoromethylphenyl)- $\alpha$ -hydroxybutyronitrile and 4-(2-hydroxyphenyl)- $\alpha$ -hydroxybutyronitrile.
4. The process as claimed in Claim 1, wherein the reaction occurs in a system having a pH of 6 to 11.
5. The process as claimed in Claim 1, wherein the yield of the compound according to formula (I) in the reaction is 50 to 100% based on the amount of the  $\alpha$ -hydroxynitrile according to formula (II).
6. The process as claimed in Claim 1, wherein the reaction occurs in the presence of a member selected from the group consisting of sodium sulfite, acid sodium sulfite, sodium dithionite, potassium sulfite, acid potassium sulfite, potassium dithionite, ammonium sulfite, acid ammonium sulfite, and ammonium dithionite, in an amount of 1 to 1000 mM.
7. The process as claimed in Claim 1, wherein the microorganism belongs to the genus Rhodococcus.
8. The process as claimed in Claim 1, wherein the microorganism belongs to the genus Alcaligenes.
9. The process as claimed in Claim 1, wherein the microorganism belongs to the genus Brevibacterium.
10. The process as claimed in Claim 1, wherein the microorganism belongs to the genus Pseudomonas.


#### Patentansprüche

1. Verfahren zur überwiegenden Herstellung einer eine Phenylgruppe aufweisenden optisch aktiven  $\alpha$ -Hydroxycarbonsäure der Formel (I):



worin X ein Wasserstoffatom, eine Hydroxylgruppe, eine Alkylgruppe mit 1 bis 3 Kohlenstoffatomen, eine Alkoxygruppe mit 1 bis 3 Kohlenstoffatomen, eine Thioalkylgruppe, ein Halogenatom, eine Phenyl-, Phenoxy-, Amino- oder eine Nitrogruppe, welche an die o-, m- oder p-Position gebunden sind, und n 1 oder 2 darstellen, wobei man einen Mikroorganismus, der dazu befähigt ist, eine Nitrilgruppe eines racemischen  $\alpha$ -Hydroxynitrils der Formel (II) asymmetrisch zu hydrolysieren:






worin X und n wie oben definiert sind,

und wobei der Mikroorganismus zum Genus *Rhodococcus*, *Alcaligenes*, *Brevibacterium* oder *Pseudomonas* oder zu behandelten Zellen dazu gehört, mit einem racemischen  $\alpha$ -Hydroxynitril der Formel (II) oder einer Mischung eines dem genannten Nitril entsprechenden Aldehyds und aus Blausäure in einem neutralen bis basischen wässrigen Medium reagieren läßt.


2. Verfahren gemäß Anspruch 1, worin der genannte Mikroorganismus die Eigenschaften von *Rhodococcus* sp. HT29-7 (FERM BP-3857), *Alcaligenes* sp. BC35-2 (FERM BP-3318), *Alcaligenes* sp. BC12-2 (FERM P-11263), *Alcaligenes* sp. BC 20 (FERM P-11264), *Alcaligenes* sp. BC24 (FERM P-12063), *Pseudomonas* sp. BC-18 (FERM BP-4536) oder von *Brevibacterium acetylicum* (IAM 1790) hat.
3. Verfahren gemäß Anspruch 1, worin das  $\alpha$ -Hydroxynitril Phenyllactonitril, 4-Phenyl- $\alpha$ -hydroxybutyronitril, 3-(2-Methoxyphenyl)lactonitril, 3-(3-Methoxyphenyl)lactonitril, 3-(4-Methoxyphenyl)lactonitril, 4-(4-Fluorphenyl)- $\alpha$ -hydroxybutyronitril, 4-(2-Chlorphenyl)- $\alpha$ -hydroxybutyronitril, 4-(4-Bromphenyl)- $\alpha$ -hydroxybutyronitril, 4-(2-Trifluormethylphenyl)- $\alpha$ -hydroxybutyronitril, 4-(3-(Trifluormethylphenyl))- $\alpha$ -hydroxybutyronitril und 4-(2-Hydroxyphenyl)- $\alpha$ -hydroxybutyronitril ist.
4. Verfahren gemäß Anspruch 1, worin die Reaktion in einem System mit einem pH-Wert von 6 bis 11 erfolgt.
5. Verfahren gemäß Anspruch 1, wobei die Ausbeute der Verbindung der Formel (I) in der Reaktion 50 bis 100% beträgt, bezogen auf die Menge des  $\alpha$ -Hydroxynitrils der Formel (II).
6. Verfahren gemäß Anspruch 1, worin die Reaktion in der Gegenwart eines Mittels, ausgewählt aus der Gruppe, bestehend aus Natriumsulfit, saurem Natriumsulfit, Natriumdithionit, Kaliumsulfit, saurem Kaliumsulfit, Kaliumdithionit, Ammoniumsulfit, saurem Ammoniumsulfit und aus Ammoniumdithionit, in einer Menge von 1 bis 1000 mM erfolgt.
7. Verfahren gemäß Anspruch 1, worin der Mikroorganismus zum Genus *Rhodococcus* gehört.
8. Verfahren gemäß Anspruch 1, worin der Mikroorganismus zum Genus *Alcaligenes* gehört.
9. Verfahren gemäß Anspruch 1, worin der Mikroorganismus zum Genus *Brevibacterium* gehört.
10. Verfahren gemäß Anspruch 1, worin der Mikroorganismus zum Genus *Pseudomonas* gehört.

#### Revendications

1. Procédé permettant de produire de manière prédominante un acide  $\alpha$ -hydroxycarboxylique optiquement actif contenant un groupe phényle, tel que représenté par la formule (I) :



dans laquelle X représente un atome d'hydrogène, un groupe hydroxyle, un groupe alkyle ayant de 1 à 3 atomes de carbone, un groupe alcoxy ayant de 1 à 3 atomes de carbone, un groupe thioalkyle, un atome d'halogène, un groupe phényle, un groupe phénoxy, un groupe amino ou un groupe nitro qui est lié en position o, m ou p ; et n vaut 1 ou 2, ledit procédé consistant à faire réagir un micro-organisme capable d'hydrolyser de manière asymétrique un groupe nitrile d'un  $\alpha$ -hydroxynitrile racémique représenté par la formule (II) :



dans laquelle X et n sont définis comme ci-dessus, et dans lequel ledit micro-organisme appartient au genre Rhodococcus, Alcaligenes, Brevibacterium ou Pseudomonas ou des cellules de ceux-ci traitées avec un  $\alpha$ -hydroxynitrile racémique représenté par la formule (II) ou bien un mélange d'un aldéhyde correspondant audit nitrile et d'acide prussique dans un milieu aqueux allant de neutre à basique.

2. Procédé selon la revendication 1, dans lequel ledit micro-organisme possède les propriétés de Rhodococcus sp. HT29-7 (FERM BP-3857), Alcaligenes sp. BC35-2 (FERM BP-3318), Alcaligenes sp. BC12-2 (FERM P-11263), Alcaligenes sp. BC20 (FERM P-11264), Alcaligenes sp. BC24 (FERM P-12063), Pseudomonas sp. BC-18 (FERM BP-4536) ou Brevibacterium acetylicum (IAM 1790).
3. Procédé selon la revendication 1, dans lequel l' $\alpha$ -hydroxynitrile est le phényllactonitrile, le 4-phényl- $\alpha$ -hydroxybutyronitrile, le 3-(2-méthoxyphényl)-lactonitrile, le 3-(3-méthoxyphényl)-lactonitrile, le 3-(4-méthoxyphényl)-lactonitrile, le 4-(4-fluorophényl)- $\alpha$ -hydroxybutyronitrile, le 4-(2-chlorophényl)- $\alpha$ -hydroxybutyronitrile, le 4-(4-bromophényl)- $\alpha$ -hydroxybutyronitrile, le 4-(2-trifluorométhylphényl)- $\alpha$ -hydroxybutyronitrile, le 4-(3-trifluorométhylphényl)- $\alpha$ -hydroxybutyronitrile et le 4-(2-hydroxyphényl)- $\alpha$ -hydroxybutyronitrile.
4. Procédé selon la revendication 1, dans lequel la réaction se fait dans un système ayant un pH de 6 à 11.
5. Procédé selon la revendication 1, dans lequel le rendement en composé suivant la formule (I) dans la réaction est de 50 à 100 % sur la base de la quantité d' $\alpha$ -hydroxynitrile suivant la formule (II).
6. Procédé selon la revendication 1, dans lequel la réaction se fait en présence d'un élément choisi dans le groupe comprenant le sulfite de sodium, le sulfite de sodium acide, le dithionite de sodium, le sulfite de potassium, le sulfite de potassium acide, le dithionite de potassium, le sulfite d'ammonium, le sulfite d'ammonium acide ou le dithionite d'ammonium, en une quantité de 1 à 1000 mM.
7. Procédé selon la revendication 1, dans lequel le micro-organisme appartient au genre Rhodococcus.
8. Procédé selon la revendication 1, dans lequel le micro-organisme appartient au genre Alcaligenes.
9. Procédé selon la revendication 1, dans lequel le micro-organisme appartient au genre Brevibacterium.
10. Procédé selon la revendication 1, dans lequel le micro-organisme appartient au genre Pseudomonas.